

4-2 Videos Guide

4-2a

- The double integral over general regions

$$\begin{aligned} & \circ \quad \iint_D f(x, y) dA \\ & = \int_a^b \int_{g_1(x)}^{g_2(x)} f(x, y) dy dx \quad \text{if } D = \{(x, y) | a \leq x \leq b, g_1(x) \leq y \leq g_2(x)\} \\ \text{OR} \quad & = \int_c^d \int_{h_1(y)}^{h_2(y)} f(x, y) dx dy \quad \text{if } D = \{(x, y) | h_1(y) \leq x \leq h_2(y), c \leq y \leq d\} \end{aligned}$$

Exercises:

4-2b

- Evaluate the iterated integral.

$$\circ \quad \int_0^2 \int_0^{y^2} x^2 y dx dy$$

4-2c

- Evaluate the double integral.
- $$\iint_D (x^2 + 2y) dA, \quad D \text{ is bounded by } y = x, y = x^3, x \geq 0$$

4-2d

- Find the volume of the solid enclosed by the paraboloid $z = x^2 + y^2 + 1$ and the planes $x = 0, y = 0, z = 0$, and $x + y = 2$.
- Sketch the region of integration and change the order of integration.

$$\int_0^2 \int_{x^2}^4 f(x, y) dy dx$$

4-2e

- Evaluate the integral by reversing the order of integration.

$$\int_0^1 \int_{x^2}^1 \sqrt{y} \sin y dy dx$$