

## 5-8 Videos Guide

### 5-8a

Theorem (statement):

- Stokes' Theorem:  $\int_C \mathbf{F} \cdot d\mathbf{r} = \iint_S \text{curl } \mathbf{F} \cdot d\mathbf{S}$ , where  $C$  is the positively oriented piecewise-smooth boundary curve of  $S$ , an oriented piecewise-smooth surface
- Green's Theorem as a special case of Stokes' Theorem via a vector form of Green's Theorem

$$\int_C \mathbf{F} \cdot d\mathbf{r} = \iint_S \text{curl } \mathbf{F} \cdot d\mathbf{S} = \iint_D (\text{curl } \mathbf{F}) \cdot \mathbf{k} \, dA$$

- Note that  $d\mathbf{S} = \mathbf{n}dS = (\mathbf{r}_u \times \mathbf{r}_v) \, dA$ , where  $\mathbf{n}$  is a unit normal vector and  $\mathbf{r}_u \times \mathbf{r}_v$  is simply a normal vector to the surface  $S$

Exercises:

### 5-8b

- Verify that Stokes' Theorem is true for the given vector field  $\mathbf{F}$  and surface  $S$ .  
 $\mathbf{F}(x, y, z) = -2yz \mathbf{i} + y \mathbf{j} + 3x \mathbf{k}$ ,  
 $S$  is the part of the paraboloid  $z = 5 - x^2 - y^2$  that lies above the plane  $z = 1$ , oriented upward

### 5-8c

- Use Stokes' Theorem to evaluate  $\iint_S \text{curl } \mathbf{F} \cdot d\mathbf{S}$ .  
 $\mathbf{F}(x, y, z) = x^2 \sin z \mathbf{i} + y^2 \mathbf{j} + xy \mathbf{k}$ ,  
 $S$  is the part of the paraboloid  $z = 1 - x^2 - y^2$  that lies above the  $xy$ -plane, oriented upward

### 5-8d

- Use Stokes' Theorem to evaluate  $\int_C \mathbf{F} \cdot d\mathbf{r}$ . In each case  $C$  is oriented counterclockwise as viewed from above.
  - $\mathbf{F}(x, y, z) = \mathbf{i} + (x + yz) \mathbf{j} + (xy - \sqrt{z}) \mathbf{k}$ ,  
 $C$  is the boundary of the part of the plane  $3x + 2y + z = 1$  in the first octant

### 5-8e

- $\mathbf{F}(x, y, z) = 2y \mathbf{i} + xz \mathbf{j} + (x + y) \mathbf{k}$ ,  
 $C$  is the curve of intersection of the plane  $z = y + 2$  and the cylinder  $x^2 + y^2 = 1$