

6.8 Videos Guide

6.8a

- l'Hôpital's Rule for indeterminate forms
 - Suppose that f and g are differentiable and $g'(x) \neq 0$ on an open interval I that contains a (except possibly at a). Suppose that

$$\lim_{x \rightarrow a} f(x) = 0 \quad \text{and} \quad \lim_{x \rightarrow a} g(x) = 0$$

or that $\lim_{x \rightarrow a} f(x) = \pm\infty$ and $\lim_{x \rightarrow a} g(x) = \pm\infty$

Then $\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)}$ if the limit on the right-hand side exists or is ∞ or $-\infty$

- Indeterminate forms
 - $\frac{0}{0}$, $\frac{\infty}{\infty}$, $0 \cdot \infty$, $\infty - \infty$, 0^0 , ∞^0 , and 1^∞

Exercises:

Find the limit. Use l'Hôpital's Rule where appropriate. If there is a more elementary method, consider using it. If l'Hôpital's Rule doesn't apply, explain why.

6.8b

- $\lim_{x \rightarrow 0} \frac{x^2}{1 - \cos x}$
- $\lim_{\theta \rightarrow \pi} \frac{1 + \cos \theta}{1 - \cos \theta}$
- $\lim_{x \rightarrow \infty} \frac{\ln \sqrt{x}}{x^2}$
- $\lim_{x \rightarrow 0} \frac{\sinh x - x}{x^3}$

6.8d

- $\lim_{x \rightarrow -\infty} x \ln \left(1 - \frac{1}{x}\right)$
- $\lim_{x \rightarrow 1^+} [\ln(x^7 - 1) - \ln(x^5 - 1)]$

6.8e

- $\lim_{x \rightarrow 0^+} (\tan 2x)^x$

6.8f

- $\lim_{x \rightarrow \infty} \left(1 + \frac{a}{x}\right)^{bx}$

6.8c

- $\lim_{t \rightarrow 0} \frac{8^t - 5^t}{t}$
- $\lim_{x \rightarrow a^+} \frac{\cos x \ln(x-a)}{\ln(e^x - e^a)}$

6.8g

Proof:

- l'Hôpital's Rule