

7.7 Videos Guide

7.7a

- The Midpoint Rule:
 - $\int_a^b f(x) dx \approx \Delta x [f(\bar{x}_1) + f(\bar{x}_2) + \cdots + f(\bar{x}_n)]$, where $\Delta x = \frac{b-a}{n}$ and $\bar{x}_i = \frac{x_{i-1} + x_i}{2}$, the midpoint of $[x_{i-1}, x_i]$
 - Error is $|E_M| \leq \frac{K(b-a)^3}{24n^2}$, where $|f''(x)| \leq K$ for $x \in [a, b]$

7.7b

- The Trapezoid Rule:
 - $\int_a^b f(x) dx \approx \frac{\Delta x}{2} [f(x_0) + 2f(x_1) + 2f(x_2) + \cdots + 2f(x_{n-1}) + f(x_n)]$, where $\Delta x = \frac{b-a}{n}$ and $x_i = a + i\Delta x$
 - Error is $|E_T| \leq \frac{K(b-a)^3}{12n^2}$, where $|f''(x)| \leq K$ for $x \in [a, b]$

Exercises:

7.7c

- Use (a) the Trapezoidal Rule and (b) the Midpoint Rule to approximate the given integral with the specified value of n . (Round your answers to six decimal places.)
 $\int_0^4 \sqrt{y} \cos y \, dy, \quad n = 8$

7.7d

- a) Estimate the errors for T_{10} and M_{10} for $\int_1^2 e^{1/x} dx$.
b) How large do we have to choose n so that the approximations T_n and M_n to the integral in part (a) are accurate to within 0.0001?

7.7e

- Simpson's Rule:
 - $\int_a^b f(x) dx \approx \frac{\Delta x}{3} [f(x_0) + 4f(x_1) + 2f(x_2) + 4f(x_3) + \cdots + 2f(x_{n-2}) + 4f(x_{n-1}) + f(x_n)]$, where n is even, $\Delta x = \frac{b-a}{n}$, and $x_i = a + i\Delta x$
 - Error is $|E_S| \leq \frac{K(b-a)^5}{180n^4}$, where $|f^{(4)}(x)| \leq K$ for $x \in [a, b]$