

16.7 Videos Guide

16.7a

- Surface integral of a scalar field $f(x, y, z)$
 - $\iint_S f(x, y, z) dS = \iint_D f(\mathbf{r}(u, v)) |\mathbf{r}_u \times \mathbf{r}_v| dA$
Note that $dS = |\mathbf{r}_u \times \mathbf{r}_v| dA$.

Exercises:

- Evaluate the surface integral.
 - $\iint_S xyz dS$,
 S is the cone with parametric equations $x = u \cos v$, $y = u \sin v$, $z = u$, $0 \leq u \leq 1$, $0 \leq v \leq \pi/2$

16.7b

- $\iint_S xy dS$,
 S is the part of the plane $2x + 2y + z = 4$ that lies in the first octant

16.7c

- If S consists of multiple surfaces S_i , then
 $\iint_S f(x, y, z) dS = \iint_{S_1} f(x, y, z) dS + \iint_{S_2} f(x, y, z) dS + \cdots + \iint_{S_n} f(x, y, z) dS$

Exercise:

- Evaluate the surface integral.
 $\iint_S (x^2 + y^2 + z^2) dS$,
 S is the part of the cylinder $x^2 + y^2 = 9$ between the planes $z = 0$ and $z = 2$, together with its top and bottom disks

16.7d

- Surface integral of a vector field $\mathbf{F}(x, y, z)$
 - Flux is $\iint_S \mathbf{F} \cdot d\mathbf{S} = \iint_S \mathbf{F} \cdot \mathbf{n} dS = \iint_D \mathbf{F} \cdot (\mathbf{r}_u \times \mathbf{r}_v) dA$
Note that $d\mathbf{S} = \mathbf{n} dS = (\mathbf{r}_u \times \mathbf{r}_v) dA$, where \mathbf{n} is a unit normal vector and $\mathbf{r}_u \times \mathbf{r}_v$ is simply a normal vector to the surface S .
 - If x and y are the parameters, we have
 $\iint_S \mathbf{F} \cdot d\mathbf{S} = \iint_D \left(-P \frac{\partial g}{\partial x} - Q \frac{\partial g}{\partial y} + R \right) dA$, for upward orientation. The signs of the integrand change for downward orientation.

Exercises:

16.7e

- Evaluate the surface integral $\iint_S \mathbf{F} \cdot d\mathbf{S}$ for the given vector field \mathbf{F} and the oriented surface S . In other words, find the flux of \mathbf{F} across S . For closed surfaces, use the positive (outward) orientation.

- $\mathbf{F}(x, y, z) = -x \mathbf{i} - y \mathbf{j} + z^3 \mathbf{k}$,
 S is the part of the cone $z = \sqrt{x^2 + y^2}$ between the planes $z = 1$ and $z = 3$ with downward orientation

16.7f

- $\mathbf{F}(x, y, z) = x \mathbf{i} + y \mathbf{j} + 5 \mathbf{k}$,
 S is the boundary of the region enclosed by the cylinder $x^2 + z^2 = 1$ and the planes $y = 0$ and $x + y = 2$