

2.3 - Linear Equations

2.3a

Definition: A first-order differential equation of the form

$a_1(x)\frac{dy}{dx} + a_0(x)y = g(x)$ is a **linear equation** in y (the dependent variable).

This is put into **standard form** by dividing both sides by $a_1(y)$ to obtain

$$\frac{dy}{dx} + P(x)y = f(x)$$

Example: Consider the DE $3\frac{dy}{dx} + 12y = 4$. In standard form, this is

$$\frac{dy}{dx} + 4y = \frac{4}{3}.$$

Where did e^{4x} come from?

2.3b

Example: Find the general solution of the given differential equation. Give the largest interval I over which the general solution is defined. Determine whether there are any transient terms in the general solution.

$$y' + 2xy = x^3$$

Example: $(1+x)\frac{dy}{dx} - xy = x + x^2$

Example: $(1+x)\frac{dy}{dx} - xy = x + x^2$



Example: $\frac{dP}{dt} + 2tP = P + 4t - 2$

[illegible]

Example: Solve the given initial-value problem. Use a graphing utility to graph the continuous function $y(x)$.

$$(1+x^2)\frac{dy}{dx} + 2xy = f(x), \quad y(0) = 0, \text{ where}$$

$$f(x) = \begin{cases} x, & 0 \leq x < 1 \\ -x & x \geq 1 \end{cases}$$

[illegible]


