

7.1 - Definition of the Laplace Transform

7.1a

Definition: A **transform** transforms one function to another function.

Examples of transforms

The derivative and the integral:

$$\frac{d}{dx}(\tan^{-1} x) = \frac{1}{1+x^2} \text{ and } \frac{d}{dx}(x^3) = 3x^2 \text{ are transforms.}$$

$$\int \frac{1}{x} dx = \ln|x| + C \text{ is a transform.}$$

Transforms possess the linearity properties, e.g.,

$$\frac{d}{dx}[af(x) + bg(x)] = a\frac{d}{dx}[f(x)] + b\frac{d}{dx}[g(x)]$$

Definition 7.1.1: Let $f(t)$ be a function defined for $t \geq 0$. Then the integral $\mathcal{L}\{f(t)\} = \int_0^\infty e^{-st} f(t) dt$ is said to be the **Laplace transform** of f , provided that the integral converges.

The Laplace transform possesses the linearity properties.

Example 1: Evaluate $\mathcal{L}\{1\}$.

Example 2: Evaluate $\mathcal{L}\{t\}$.

Example: Evaluate $\mathcal{L}\{4t - 7\}$

7.1b

Example: Evaluate $\mathcal{L}\{\sin kt\}$

Theorem 7.1.1: Transforms of Some Basic Functions

$$(a) \mathcal{L}\{1\} = \frac{1}{s}$$

$$(b) \mathcal{L}\{t^n\} = \frac{n!}{s^{n+1}}, n = 1, 2, 3, \dots$$

$$(c) \mathcal{L}\{e^{at}\} = \frac{1}{s-a}$$

$$(d) \mathcal{L}\{\sin kt\} = \frac{k}{s^2 + k^2}$$

$$(e) \mathcal{L}\{\cos kt\} = \frac{s}{s^2 + k^2}$$

$$(f) \mathcal{L}\{\sinh kt\} = \frac{k}{s^2 - k^2}$$

$$(g) \mathcal{L}\{\cosh kt\} = \frac{s}{s^2 - k^2}$$

7.1c

Definition: A function f is said to be of **exponential order** if there exist constants $c, M > 0$, and $T > 0$ such that $|f(t)| \leq Me^{ct}$ for all $t > T$.

Theorem: Sufficient Conditions for Existence

[illegible]

Theorem: Behavior of $F(s)$ as $s \rightarrow \infty$

If f is piecewise continuous on $[0, \infty)$ and of exponential order and $F(s) = \mathcal{L}\{f(t)\}$, then $\lim_{s \rightarrow \infty} F(s) = 0$.

Example: Use Definition 7.1.1 to find $\mathcal{L}\{f(t)\}$.

7.1d

$$f(t) = \begin{cases} 2t + 1, & 0 \leq t < 1 \\ 0, & t \geq 1 \end{cases}$$

Example: Use Definition 7.1.1 to find $\mathcal{L}\{f(t)\}$.

$$f(t) = e^{-2t-5}$$

Example: Use Theorem 7.1.1 to find $\mathcal{L}\{f(t)\}$.

$$f(t) = t^2 - e^{-9t} + 5$$

$$f(t) = \cos^2 t$$
