6.4 - Special Functions

6.4a

Bessel's equation of order ν

$x^2y'' + xy' + (x^2 - \nu^2)y = 0$

Example: Find the general solution of the given differential equation on $(0, \infty)$.

$16x^2y'' + 16xy' + (16x^2 - 1)y = 0$

Bessel Functions of the First and Second Kinds

Assuming a solution of the form $y = \sum_{n=0}^{\infty} c_n x^{n+r}$ and substituting into Bessel's equation, we have

$$\sum_{n=0}^{\infty}(n+r)(n+r-1)c_n x^{n+r} + \sum_{n=0}^{\infty}(n+r)c_n x^{n+r} + \sum_{n=0}^{\infty} c_n x^{n+r+2} - \sum_{n=0}^{\infty} \nu^2 c_n x^{n+r} = 0$$

$$\implies \sum_{n=0}^{\infty}[(n+r)(n+r) - \nu^2]c_n x^{n+r} + \sum_{n=0}^{\infty} c_n x^{n+r+2} = 0$$

Reindexing yields $\sum_{k=0}^{\infty}[(k+r)^2 - \nu^2]c_k x^{k+r} + \sum_{k=2}^{\infty} c_{k-2} x^{k+r} = 0$

For $k = 0$ we find $r = \pm\nu$ and $c_1 = 0$.

Then $c_k = -\dfrac{c_{k-2}}{(k+r)^2 - \nu^2}$, $k = 2,\ 3,\ 4,\ \ldots$

A standard choice for c_0 is $c_0 = \dfrac{1}{2^\nu \Gamma(1+\nu)}$, where $\Gamma(1+\nu)$ is the gamma function.

Definition: The **gamma function** is the integral-defined function

$$\Gamma(x) = \int_0^\infty t^{x-1} e^{-t} dt$$

The function $y = x^\nu \sum_{n=0}^{\infty} c_{2n} x^{2n}$ is then

$$y = \sum_{n=0}^{\infty} \frac{(-1)^n}{n!\Gamma(1+\nu+n)} \left(\frac{x}{2}\right)^{2n+\nu} = J_\nu(x)$$

We also have that $J_{-\nu}(x) = \sum_{n=0}^{\infty} \frac{(-1)^n}{n!\Gamma(1-\nu+n)} \left(\frac{x}{2}\right)^{2n-\nu}$

6.4b

Definition: The functions $J_\nu(x)$ and $J_{-\nu}(x)$ defined above are **Bessel functions of the first kind** of order ν and $-\nu$, respectively, and converge at least on $(0, \infty)$.

If ν is not an integer, then the general solution to Bessel's equation is $y = c_1 J_\nu(x) + c_2 J_{-\nu}(x)$.

Definition: The function $Y_\nu(x) = \frac{\cos(\nu\pi) J_\nu(x) - J_{-\nu}(x)}{\sin \nu\pi}$ (ν a non-integer) is a **Bessel function of the second kind**. It can be shown to be linearly independent with $J_\nu(x)$.

If ν is an integer, then it can be shown that $J_\nu(x)$ and $J_{-\nu}(x)$ are linearly dependent. However, $J_\nu(x)$ and $Y_\nu(x)$ are independent. As such, for an integer value of ν, we use $J_\nu(x)$ and $Y_\nu(x)$ (in this case, $Y_\nu(x)$ is actually defined by a limiting process as $\nu \to m$, where m is an integer). In this case we can give the general solution to Bessel's equation as $y = c_1 J_\nu(x) + c_2 Y_\nu(x)$.

Example: Find the general solution of the given differential equation on $(0, \infty)$.

$x^2 y'' + xy' + (x^2 - 1)y = 0$

Alternate forms of Bessel's equation

Using a substitution we can find that the general solution to the DE $x^2y'' + xy' + (\alpha^2x^2 - \nu^2)y = 0$ is $y = c_1J_\nu(\alpha x) + c_2Y_\nu(\alpha x)$.

Definition: The preceding differential equation is called the **parametric Bessel equation of order ν**.

Definition: The differential equation $x^2y'' + xy' - (x^2 + \nu^2)y = 0$ is the **modified Bessel equation of order ν**. It can be converted to a Bessel equation by the substitution $t = ix$, where $i^2 = -1$; the result is...

Definition: The **modified Bessel function of the first kind** of order ν, $I_\nu(x) = i^{-\nu}J_\nu(ix)$.

The general solution to the modified Bessel equation of order ν (ν not an integer) is $y = c_1I_\nu(x) + c_2I_{-\nu}(x)$.

Definition: As before, for non-integer values of ν we define the **modified Bessel function of the second kind**, $K_\nu(x) = \dfrac{\pi}{2}\dfrac{I_{-\nu}(x) - I_\nu(x)}{\sin\nu\pi}$ and extend this definition to integer values using a limit process similar to that referenced when defining $Y_\nu(x)$.

And as before, the general solution to the modified Bessel equation of order ν (for ν an integer) is $y = c_1I_\nu(x) + c_2K_\nu(x)$. A parametric form of the modified Bessel equation and associated solution exists such as we saw above.

Definition: An equation of the form $x^2y'' + xy' - (\alpha^2x^2 + \nu^2)y = 0$ is a **parametric form of the modified Bessel equation of order ν** and can be obtained from a modified Bessel equation using a change of variables.

The general solution for such an equation is $y = c_1I_\nu(\alpha x) + c_2K_\nu(\alpha x)$.

Example: Find the general solution of the given differential equation on $(0, \infty)$.

$x^2y'' + xy' - (2x^2 + 64)y = 0$

Example: The general solution to

$y''+\frac{1-2a}{x}y'+\left(b^2c^2x^{2c-2}+\frac{a^2-p^2c^2}{x^2}\right)y=0, p \geq 0$ is $y = x^a\left[c_1J_p\left(bx^c\right)+c_2Y_p\left(bx^c\right.\right.$

Use this to find the general solution of the given differential equation on $(0, \infty)$.

$xy'' + 3y' + xy = 0$

Legendre polynomials

Definition: The differential equation

$$(1 - x^2)y'' - 2xy' + n(n + 1)y = 0$$

is **Legendre's equation of order *n***.

Note that $x = 0$ is an ordinary point. Starting with $y = \sum_{k=0}^{\infty} c_k x^k$,...

leads to $2c_2 + 6c_3 x - 2c_1 x + n(n+1)c_0 + n(n+1)c_1 x$
$+ \sum_{j=2}^{\infty} \left\{(j+2)(j+1)c_{j+2} + [-j(j-1) - 2j + n(n+1)]\, c_j\right\} x^j = 0$

We find that $c_{j+2} = -\dfrac{(n-j)(n+j+1)}{(j+2)(j+1)} c_j$, $j = 2,\ 3,\ 4,\ \ldots$

$$c_2 = -\frac{n(n+1)}{2} c_0$$
$$c_4 = -\frac{(n-2)(n+3)}{4 \cdot 3} c_2 = \frac{(n-2)n(n+1)(n+3)}{4!} c_0$$

This terminates if n is even.

$$c_3 = -\frac{(n-1)(n+2)}{6} c_1$$
$$c_5 = -\frac{(n+4)(n-3)}{5 \cdot 4} c_3 = \frac{(n-3)(n-1)(n+2)(n+4)}{5!} c_1$$

This terminates if n is odd.

Traditional values for c_i are

For $n = 0$, $c_0 = 1$; for $n = 2,\ 4,\ 6,\ \ldots$, $c_0 = (-1)^{n/2} \dfrac{1 \cdot 3 \cdots (n-1)}{2 \cdot 4 \cdots n}$

For $n = 1$, $c_1 = 1$; for $n = 3,\ 5,\ 7,\ \ldots$, $c_1 = (-1)^{(n-1)/2} \dfrac{1 \cdot 3 \cdots n}{2 \cdot 4 \cdots (n-1)}$

Definition: For each value of n the result is a **Legendre polynomial of order *n***.

The first six Legendre polynomials are

$$P_0(x) = 1 \qquad P_1(x) = x$$
$$P_2(x) = \frac{1}{2}(3x^2 - 1) \qquad P_3(x) = \frac{1}{2}(5x^3 - 3x)$$
$$P_4(x) = \frac{1}{8}(35x^4 - 30x^2 + 3) \qquad P_5(x) = \frac{1}{8}(63x^5 - 70x^3 + 15x)$$